arXiv:physics/0310091v1 [physics.optics] 18 Oct 2003

Comments on "Geometric Phase Associated with Transformations of Optical Beams Bearing Orbital Angular Momentum"

S.C. Tiwari

Institute of Natural Philosophy,

1, Kusum Kutir, Mahamanapuri,

Varanasi - 221 005

The first direct measurement of the geometric phase acquired by the light beams carrying orbital angular momentum (OAM) has been reported by Galvez et al [1]. The closed path traversed by the optical beams is the space of modes of Laguerre-Gaussian beams [2]. Modes LG_p^l carry OAM of $l\hbar$ per photon, where l is the azimuthal index, and p is the radial index. In the experiments, a geometric phase on a LG_0^{-1} beam is introduced, changed, and interfered with a reference LG_0^0 beam. Authors conclude the paper with the statements: "All of these geometric phases arose from transformations involving a change in the OAM of the modes. They support the conjecture that geometric phase arises from the exchange of angular momentum between the light and optical system [13]". Ref.13 in [1] is an interesting work by van Enk,here after referred at [3]. In the present note I make following comments:

A1. The connection of angular momentum to the geometrical phase in optics was vaguely hinted at in [4]. In 1992 plausible arguments were presented to propose that the physical mechanism of geometrical phase in optics was the transfer of angular momentum between the light and optical elements [5]. This conjecture was theoretically supported by van Enk [3] ,and later by Banerjee [6]. In [3] van Enk considered LG beams for the geometric phases.

A2. In 1994, LeFloch showed interest to verify this conjecture in his laboratory, however the extreme difficulty in the measurement of angular momentum transfer was pointed out by him [7]. The present work measures geometric phase, and is significant; however the direct demonstration of angular momentum exchange remains elusive. I believe a direct experimental verification of this relationship, and measurement of angular momentum exchange would have far reaching implications on the nature of light and crystal optics [8].

Acknowledgement

The library facility of Banaras Hindu University, Varanasi is acknowledged.

---

[1] E.J. Galvez et al, Phys. Rev. Lett. **90**, 203901(2003)

[2] L. Allen et al,Phys. Rev. A **45**, 8185(1992)

[3] S.J. van Enk, Opt.Commun. **102**, 59(1993)

[4] H.Jiao et al, Phys. Rev. A **39**,3475(1989); S.C. Tiwari, Phys. Lett. A **149**, 223(1990)

[5] S.C. Tiwari, J. Mod. Optics **39**, 1097(1992)

[6] D. Banerjee, Phys. Rev. E **56**, 1129(1997)

[7] A. LeFloch, Private Communication(1994)

[8] S.C. Tiwari, J. Optics B, **4**, S39(2002)